Parameter Values for Reactor Problem

$$\begin{array}{llll} V & = & 0.01 & m^3 \\ C_{A0} & = & 1 & kgmol/m^3 \\ \rho & = & 850 & kg/m^3 \\ C_p & = & 1 & kJ/kg/K \\ (-\Delta H_1) & = & 4.00 \text{x} 10^4 & kJ/kgmole \\ (-\Delta H_2) & = & 8.50 \text{x} 10^4 & kJ/kgmole \\ (-\Delta H_3) & = & 6.00 \text{x} 10^4 & kJ/kgmole \\ T_0 & = & 200 & {}^{\circ}C \\ k_{10} & = & 100 & s^{-1} \\ k_{20} & = & 150 & s^{-1} \\ k_{30} & = & 1.1 & m^3/(kgmole)/s \\ E_1 & = & 3.00 \text{x} 10^4 & kJ/Kgmole/K \\ E_2 & = & 4.00 \text{x} 10^4 & kJ/Kgmole/K \\ E_3 & = & 2.00 \text{x} 10^4 & kJ/Kgmole/K \end{array}$$

• The rate constants, k_i are related to temperature by:

$$k_i = k_{i0} exp\left(-\frac{E_i}{RT}\right)$$

• There are 3 reactions occurring in the CSTR. The associated heats of reaction are given in the table below:

	reaction	heat of reaction
1	$A \to B$	$(-\Delta H_1)$
2	$B \to C$	$(-\Delta H_2)$
3	$2A \to D$	$(-\Delta H_3)$

• The reactions are exothermic as you can tell from the values of the heats of reaction. Thus Q will be a negative value.

Steady State Values of Inputs

$$F_s = 7x10^{-5} m^3/s$$

 $Q_s = -7 kJ/s$