


Figure 5.3. Particle-size distribution curves for sediments in Czechoslovakia (Bazant, 1979): 1. Vltava River gravel; 2. "Gap-graded" gravel; 3. Letna terrace, uniform sand; 4. Pankvac terrace, gap-graded clayey sand; 5. Micovna loess; 6. Hodonin silt; 7. Ruzyne clay; 8. Branany bentonite; 9. Quartzite talus from Boulder Mountain, Black Hills, South Dakota.

Casagrande Chart



UNIFIED SOIL CLASSIFICATION
(Including Identification and Description)

Major Divisions		Group Symbols	Typical Names	Field Identification Procedures (Excluding particles larger than 3 in. and basing fractions on estimated weights)			Information Required for Describing Soils		
1	2	3	4	5			6		
<p align="center">Coarse-grained Soils</p> <p align="center">More than half of material is larger than No. 200 sieve size.</p>	<p align="center">Gravels</p> <p align="center">More than half of coarse fraction is larger than No. 4 sieve size.</p> <p align="center">(For visual classification, the 1/2-in. size may be used as equivalent to the No. 4 sieve size.)</p>	GW	Well-graded gravels, gravel-sand mixture, little or no fines.	Wide range in grain size and substantial amounts of all intermediate particle sizes.			<p>For undisturbed soils add information on stratification, degree of compactness, cementation, moisture condition, and drainage characteristics.</p> <p>Give typical name; indicate approximate percentages of sand and gravel, maximum size; angularity, surface condition, and hardness of the coarse grains; local or geologic name and other pertinent descriptive information; and symbol in parentheses.</p> <p>Example: Silty sand, gravelly; about 20% hard, angular gravel particles 1/2-in. maximum size; rounded and subangular sand grains, coarse to fine; about 15% nonplastic fines with low dry strength; well compacted and moist in place; alluvial sand; (SM).</p>		
		GP	Poorly graded gravels or gravel-sand mixture, little or no fines.	Predominantly one size or a range of sizes with some intermediate sizes missing.					
		GM	Silty gravels, gravel-and-silt mixtures.	Nonplastic fines or fines with low plasticity (for identification procedures see ML below).					
		GC	Clayey gravels, gravel-and-clay mixtures.	Plastic fines (for identification procedures see CL below).					
	<p align="center">Sands</p> <p align="center">More than half of coarse fraction is smaller than No. 4 sieve size.</p> <p align="center">(For visual classification, the 1/2-in. size may be used as equivalent to the No. 4 sieve size.)</p>	<p align="center">Clean Gravels (Little or no fines)</p>	SW	Well-graded sands, gravelly sands, little or no fines.	Wide range in grain size and substantial amounts of all intermediate particle sizes.				
			SP	Poorly graded sands or gravelly sands, little or no fines.	Predominantly one size or a range of sizes with some intermediate sizes missing.				
		<p align="center">Sands with Fines (Appreciable amount of fines)</p>	SM	Silty sands, sand-silt mixtures.	Nonplastic fines or fines with low plasticity (for identification procedures see ML below).				
			SC	Clayey sands, sand-clay mixtures.	Plastic fines (for identification procedures see CL below).				
						Identification Procedure on Fraction Smaller than No. 40 Sieve Size.			
						Dry Strength (Crushing Characteristics)		Dilatancy (Reaction to shaking)	Toughness (Consistency near PL)
<p align="center">Fine-grained Soils</p> <p align="center">More than half of material is smaller than No. 200 sieve size.</p>	<p align="center">Silts and Clays Liquid Limit is less than 50</p>	ML	Inorganic silts and very fine sands, rock flour, silty or clayey fine sands or clayey silts with slight plasticity	None to slight	Quick to slow	None	<p>For undisturbed soils add information on structure, stratification, consistency in undisturbed and remolded states, moisture and drainage conditions</p>		
		CL	Inorganic clays of low to medium plasticity, gravelly clays, sandy clays, silty clays, lean clays.	Medium to high	None to very slow	Medium			
	<p align="center">Silts and Clays Liquid Limit is greater than 50</p>	OL	Organic silts and organic silty clays of low plasticity.	Slight to medium	Slow	Slight	<p>Give typical name; indicate degree and character of plasticity; amount and maximum size of coarse grains; color in wet condition; odor, if any; local or geologic name and other pertinent descriptive information; and symbol in parentheses.</p> <p>Example: Clayey silt, brown; slightly plastic; small percentage of fine sand; numerous vertical root holes; firm and dry in place; loess; (ML)</p>		
		MH	Inorganic silts, micaceous or diatomaceous fine sandy or silty soils, elastic silts.	Slight to medium	Slow to none	Slight to medium			
		CH	Inorganic clays of high plasticity, fat clays.	High to very high	None	High			
		OH	Organic clays of medium to high plasticity, organic silts.	Medium to high	None to very slow	Slight to medium			
		Pt	Peat and other highly organic soils.	Readily identified by color, odor, spongy feel and frequently by fibrous texture					

Table 4.1 AASHTO Soil Classification System

General classification	Granular materials (35% or less passing US No. 200 sieve)			Silt-clay materials (More than 35% passing US No. 200 sieve)							
	A-1		A-3	A-2				A-4	A-5	A-6	A-7
Group classification	A-1a	A-1b		A-2-4	A-2-5	A-2-6	A-2-7				A-7-5 A-7-6
Sieve analysis											
Percent passing											
US No. 10 (2 mm)	50 max										
US No. 40 (420 μ)	30 max	50 max	51 max								
US No. 200 (75 μ)	15 max	25 max	10 max	35 max	35 max	35 max	35 max	36 min	36 min	36 min	36 min
Characteristics of fraction passing US No. 40 (420 μ)											
Liquid limit											
Plasticity index											
	6 max		Non-plastic	40 max 10 max	41 min 10 max	40 max 11 min	41 min 11 min	40 max 10 max	41 min 10 max	40 max 11 min	41 min 11 min
Group index	0		0	0		4 max		8 max	12 max	16 max	20 max
Usual types of significant constituent materials	Stone fragments gravel and sand		Fine Sand	Silty or clayey gravel and sand				Silty soils		Clayey soils	
General rating as subgrade	Excellent to good							Fair to poor			

Note: A-8 is identified by visual classification, and is not shown in the Table.

Classification procedure: Proceeding from left to right in the chart, the correct group will be found by the process of elimination. The first group from the left consistent with the test data is the correct classification. A-7 group is subdivided into A-7-5 or A-7-6 depending on the plastic limit. For $w_p < 30$, the classification is A-7-6; for $w_p \geq 30$, it is A-7-5.

